

Short Malnourished Children and Fat Accumulation With Food Supplementation

Christian Fabiansen, MD, PhD,^{a,b} Kevin P.Q. Phelan, MSc,^c Bernardette Cichon, PhD,^{a,b} Charles W. Yaméogo, PhD,^{a,d} Ann-Sophie Iuel-Brockdorff, PhD,^{a,b} Anura Kurpad, MD, PhD,^e Jonathan C. Wells, PhD,^f Christian Ritz, PhD,^a Suzanne Filteau, PhD,^g André Briend, MD, PhD,^{a,h} Vibeke B. Christensen, MD, DMSc,^{a,i} Kim F. Michaelsen, MD, DMSc,^a Susan Shepherd, MD,^c Henrik Friis, MD, PhD^a

abstract

BACKGROUND: In moderate acute malnutrition programs, it is common practice to not measure mid-upper arm circumference (MUAC) of children whose length is <67 cm. This is based on expert opinion that supplementation of shorter children with low MUAC and weight-for-height z score ≥ -2 may increase risk of excessive fat accumulation. Our aim was to assess if shorter children gain more fat than taller children when treated for moderate acute malnutrition diagnosed by low MUAC alone.

METHODS: In this observational study, we included children aged 6 to 23 months with a MUAC between 115 and 125 mm and a weight-for-height z score ≥ -2 . On the basis of length at admission, children were categorized as short if <67 cm and long if ≥ 67 cm. Linear mixed-effects models were used to assess body composition on the basis of deuterium dilution and skinfold thickness.

RESULTS: After 12 weeks of supplementation, there was no difference in change in fat mass index (-0.038 kg/m², 95% confidence interval [CI]: -0.257 to 0.181 , $P = .74$) or fat-free mass index (0.061 kg/m², 95% CI: -0.150 to 0.271 , $P = .57$) in short versus long. In absolute terms, the short children gained both less fat-free mass (-230 g, 95% CI: -355 to -106 , $P < .001$) and fat mass (-97 g, 95% CI: -205 to 10 , $P = .076$). There was no difference in changes in absolute subscapular and triceps skinfold thickness and z scores ($P > .5$).

CONCLUSIONS: Short children with low MUAC do not gain excessive fat during supplementation. With these data, we support a recommendation for policy change to include all children ≥ 6 months with low MUAC in supplementary feeding programs, regardless of length. The use of length as a criterion for measuring MUAC to determine treatment eligibility should be discontinued in policy and practice.

^aDepartment of Nutrition, Exercise and Sports, University of Copenhagen, Copenhagen, Denmark; ^bMédecins Sans Frontières, Denmark, Copenhagen, Denmark; ^cAlliance for International Medical Action, Dakar, Sénégal; ^dDépartement Biomédical et Santé Publique, Institut de Recherche en Sciences de la Santé, Ouagadougou, Burkina Faso; ^eDivision of Nutrition, St John's Research Institute, Koramangala, Bangalore, India; ^fChildhood Nutrition Research Centre, University College London Great Ormond Street Institute of Child Health, London, United Kingdom; ^gDepartment of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, United Kingdom; ^hTampere Center for Child Health Research, School of Medicine, University of Tampere and Tampere University Hospital, Tampere, Finland; and ⁱDepartment of Paediatrics, Rigshospitalet, Copenhagen, Denmark

Dr Fabiansen conceptualized the research, conducted the research, acquired the data, performed statistical analysis, and wrote the first draft of the manuscript; Mr Phelan and Drs Yaméogo, Kurpad, Wells, Filteau, Briend, Christensen, Cichon, and Iuel-Brockdorff conducted the research and acquired the data; Dr Ritz conceptualized the research and design and performed statistical analysis and interpretation of data; Dr Shepherd conceptualized the research and contributed

WHAT'S KNOWN ON THIS SUBJECT: Policy for national malnutrition programs in many African countries is to exclude shorter children with moderate acute malnutrition from treatment because of concerns they will accumulate fat and be at later risk of noncommunicable diseases.

WHAT THIS STUDY ADDS: We show that shorter children with low mid-upper arm circumference do not gain excessive fat during supplementation and thus should be offered treatment. This evidence should be integrated in guidelines by the World Health Organization.

To cite: Fabiansen C, Phelan K.P.Q., Cichon B, et al. Short Malnourished Children and Fat Accumulation With Food Supplementation. *Pediatrics*. 2018;142(3):e20180679

Childhood malnutrition contributes to almost half the mortality in children <5 years.¹ Moderate acute malnutrition (MAM) is currently defined as a weight-for-height z score (WHZ) between -3 and -2 (ie, moderate wasting) and/or a mid-upper arm circumference (MUAC) between 115 and 125 mm.² Although the prevalence of MAM is unknown, moderate wasting alone affects 33 million children at any time³ and is associated with a threefold increased risk of death.⁴

Protocols for management of acute malnutrition in many African countries (including Cameroon, Central African Republic, Chad, Guinea, Ivory Coast, Mali, Mauritania, Senegal, and Togo) instruct health personnel to measure the MUAC only of children with a length ≥ 67 cm when assessing eligibility for MAM or severe acute malnutrition (SAM) treatment programs.⁵⁻¹³ In Ethiopia, admission by MUAC alone for SAM treatment is restricted to children with lengths >65 cm.¹⁴ These shorter children are enrolled in treatment only if they meet WHZ criterion, which is less closely linked to the risk of death and misses children at high risk who would have been identified by MUAC. Hence, current practice excludes short children who would benefit from nutritional support.

The practice of restricting treatment admission by MUAC alone to children of lengths ≥ 67 cm is not supported by data and has never been formally endorsed by the World Health Organization (WHO). Two reasons can be identified to explain this exclusion. First, the inclusion of a length restriction for MUAC is likely a holdover from older versions of emergency nutrition guidelines that used length as a proxy for age (<6 months) because it can sometimes be difficult to ascertain age in a fast-moving emergency. WHO, for example, used a length of 60 cm as a proxy for an

age of 6 months in publications from 1995 to 2000,^{15,16} whereas a length of 65 cm was used in older versions of guidelines from nongovernmental humanitarian organizations like Médecins Sans Frontières in 1995 and Valid International in 2006.^{17,18} It is important to identify children <6 months of age because they should be exclusively breastfed and treated as inpatients if in need of treatment. However, length is a poor proxy for age, and most often a caretaker can recall the month of birth for an infant.

Second, WHO now recommends against routine food supplementation of children with MAM because of a concern it may promote obesity and increase the risk of noncommunicable diseases later in life.¹⁹ Some experts worry that shorter children >6 months of age who only meet the low-MUAC criteria for MAM are most at risk for excessive fat accumulation during treatment.² Although only based on expert opinion, this seems to be the reason why many national malnutrition protocols exclude shorter children from having their MUAC measured, thereby making children ineligible for treatment, unless they meet WHZ criteria. This concern led WHO to call for research to identify appropriate MUAC admission and discharge criteria for children <67 cm and ≥ 6 months for supplementary feeding programs.²

We have previously shown that when supplemented, ponderal growth rates are similar in short and long children ≥ 6 months with a low MUAC.²⁰ Niger dropped this length restriction in 2016, and its revised protocols now call for measuring MUAC in all children ≥ 6 months, regardless of length.²¹

The extent to which there is a difference in fat accumulation during treatment has not been assessed. Our objective in this article is to assess if short children (<67 cm) gain more fat than long children (≥ 67 cm) when

treated for MAM diagnosed by low MUAC (115–125 mm).

METHODS

As previously reported,²⁰ a prospective cohort study was nested in a randomized nutrition trial (Treatfood)²² in which we investigated the effectiveness of 500 kcal/day supplements as either corn-soy blend (CSB) porridge or ready-to-use lipid-based nutrient supplements (LNSs). The LNSs provided almost 3 times more energy as fat than the CSB supplements (~57% vs ~21%). Assignment to 1 of the 12 supplements (6 CSBs and 6 LNSs) followed randomization stratified by site. The trial was approved by the Ethics Committee for Health Research in Burkina Faso (2012-8-059), and consultative approval was obtained from the Danish National Committee on Biomedical Research Ethics (1208204). Consent was obtained from caregivers, before inclusion, verbally and in writing (signature or fingerprints). All children were treated free of charge at the study sites, irrespective of study participation.

Participants

Data were collected in the Province du Passoré in the northern region of Burkina Faso at 5 research sites located at different governmental health centers and staffed by the nongovernmental organization Alliance for International Medical Action (Dakar, Senegal).

Children were screened in villages either by community health workers using MUAC tapes or by designated screening teams with the use of both MUAC and WHZ. Moreover, children could be referred from a health center or could present at site on a caretaker's initiative. At the sites, the final assessment of eligibility for inclusion was performed.

For the main Treatfood trial, children aged 6 to 23 months with MAM (defined as a MUAC between 115 and 124 mm and/or a WHZ between -3 and -2), who were residents in the catchment area and whose caretaker consented to participate, were included. Children were not included if they had been treated for SAM or hospitalized within the past 2 months, if they were already in a nutritional program, or if they presented medical complications requiring hospitalization. Likewise, children with a severe disability limiting the possibility of investigations and children with a suspected allergy to milk, peanuts, corn, or soy were not included.

Of the 1609 children included in the trial, 468 had a MUAC between 115 and 125 mm and a WHZ ≥ -2 and were included in the current study. In many settings, these children are excluded from treatment if they are also short (ie, <65 or 67 cm). Accordingly, children in this study were categorized, on the basis of baseline length, as short (<67 cm) or long (≥ 67 cm) to assess if short children gain more fat than long children when treated for MAM diagnosed by low MUAC.

Procedures and Study Visits

We previously described clinic visits, standard anthropometric measurements, and age determination.²⁰ In the present article, we additionally report on indices of body composition and skinfolds assessed at baseline and at the end of intervention at 12 weeks. Total body water (TBW) was assessed by using the deuterium dilution technique to enable calculation of fat-free mass (FFM) and fat mass (FM). The method involved giving an oral dose of 5 g of deuterium oxide (D_2O) (99.8%; Cambridge Isotope Laboratories

Inc, Andover, MA). The isotope was diluted in 5 g of bottled water (LAFI, Burkina Faso), with the dosing bottle weighed with 0.01 g precision (model CQT 202; Adam Equipment, United Kingdom) before and after administration of the dose. Predose saliva samples were obtained to assess background isotope levels in body fluids, and postdose saliva samples were collected after a 3-hour equilibration period as established during the pilot study.²³ For each assessment, D_2O enrichment was measured in duplicate in the pre- and postdose saliva samples and in a diluted sample of the dose by using Fourier Transform Infrared Spectrometry (Agilent Technologies, Santa Clara, CA)²⁴ at St. John's Research Centre, Bangalore, India. Saliva samples required at least 60 μL of saliva for analysis. Deuterium dilution space was calculated as described previously²⁵ and converted to TBW by using a factor of 1.044 to adjust for proton exchange.²⁶ FFM was calculated as TBW divided by hydration by using age- and sex-specific hydration coefficients.²⁷ FM was calculated as weight minus FFM. Data were cleaned for typographical errors and implausible TBW values on the basis of the association of TBW with length and a cutoff for FM of < -0.1 and > 2.4 kg.

Skinfold thickness was measured in duplicate by a Harpenden caliper. The mean of the duplicate measurements was taken for analysis. Weight was measured in duplicate to the nearest 100 g by using electronic scales (Seca model 881 1021659) with double weighing function. Length was measured in duplicate with a wooden length board to the nearest millimeter. WHZ was determined at sites by using WHO field tables, and this value was used for recruitment. MUAC was measured in duplicate to the nearest millimeter, at the midpoint between

the olecranon and the acromion process by using a standard measuring tape. Anthropometric measurements were undertaken by trained staff after standardization sessions.

In later analyses, WHZ, length-for-age z score (LAZ), and weight-for-age z score were calculated by using the package "z score 06" in Stata 12 (Stata Corp, College Station, TX). Skinfold-for-age z scores were calculated by using WHO's Anthro Plus software (version 3.2.2, 2011; WHO, Geneva, Switzerland). All z scores were calculated by using the 2006 WHO child growth standards.

Outcomes

Changes were evaluated in FM, FFM, weight, length, and skinfold thickness (both raw values and skinfold thickness-for-age z scores). By dividing FFM and FM by length squared, the indices are expressed as kg/m^2 independent of length, that is, fat-free mass index (FFMI) and fat mass index (FMI). In addition, fat was calculated as a percentage of total body weight.

Statistical Analysis

Data were double entered in Epidata 3.1 (Epidata Association, Odense, Denmark), and double entry checks were performed on a daily basis. All statistical analyses were conducted by using the statistical software package Stata version 12 (Stata Corp). Baseline and endline characteristics of children were summarized as mean (SD) or percentages, and categories were compared at baseline by using 2-sample *t* tests for continuous variables and χ^2 tests for categorical variables.

To evaluate differences in outcomes between groups during the 12-week supplementation intervention, a

linear mixed analysis of covariance model was considered the main analysis, resulting in an estimated mean difference at the end of the study adjusted for differences at baseline. Specifically, in the analyses, we included adjustment for food intervention as well as a number of baseline characteristics: baseline measure of the outcome, age, sex, and month of admission (as fixed effects). Random intercepts were included in the model to adjust for variation between sites. In the adjusted analysis, weight was derived by adding FFM and FM. Additionally, for direct outcomes of fat (FM and FMI), we evaluated the 2-way interaction between length group and product group (CSB compared with LNS) to assess effect modification by type of product. Moreover, possible effect modification was also evaluated for stunting at admission in (1) 2 categories: greater than and less than a LAZ of <-2 , or (2) 3 categories: severe stunting (LAZ <-3), moderate stunting (LAZ ≥-3 to <-2), or absence of stunting (LAZ ≥-2). The same adjustments were applied in these additional analyses as in the main analysis. Model checking was based on residuals and normal probability plots. A significance level of 0.05 was applied.

RESULTS

A total of 1609 children, of which 55% were girls, were randomly assigned to 12 weeks of supplementary feeding in the Treatfood trial.²² Of these, 50% (804) were included by both WHZ and MUAC, 21% (337) were included by WHZ alone, and 29% (468) were included by MUAC alone. Among the 468 children recruited by MUAC only, on whom this article is based, 230 (49%) were <67 cm (short), and 238 (51%) were ≥ 67 cm (long).²⁰ During the intervention, 3 short and




FIGURE 1

Participant flowchart. Treatfood participants were included in the cohort study. Adapted from Fabiansen C, Yaméogo CW, Iuel-Brockdorf A-S, et al. Effectiveness of food supplements in increasing fat-free tissue accretion in children with moderate acute malnutrition: a randomised $2 \times 2 \times 3$ factorial trial in Burkina Faso. *PLoS Med.* 2017;14(9):e1002387. ^a Body composition assessment by D_2O .

11 long children dropped out before completion of supplementation at week 12. Body composition based on D_2O dilution both at baseline and after 12 weeks were successfully determined in 195 (85%) of short children and 195 (82%) of long children (Fig 1). At baseline, there were substantial differences in age and sex distributions as well as in standard anthropometry between short and long, whereas there were no differences in allocation to the different experimental diets and season of inclusion (Table 1). Mean values of body composition variables determined at baseline and after supplementation are presented in Table 2. At baseline, the weight of short children was 1.474 (95% confidence interval [CI]: 1.355 to 1.593) kg less than that of long children, which reflected a 0.120 (95% CI: 0.047 to 0.193) kg lower FM and a 1.354 (95% CI: 1.219 to 1.489) kg lower FFM. Adjusted for length squared, this corresponded to a 0.325 (95% CI: 0.163 to 0.487) kg/m^2 higher FMI in short compared with long children, but no difference ($-0.056 kg/m^2$; 95% CI: -0.230 to 0.118) in FFMI. Body fat was 1.9

(95% CI: 0.8 to 3.0) percentage points higher in short compared with long children.

Adjusted differences from baseline to endline are presented in Table 3. After the 12 weeks of supplementation, there was no difference of change in FMI ($-0.038 kg/m^2$, 95% CI: -0.257 to 0.181, $P = .74$) or FFMI ($0.061 kg/m^2$, 95% CI: -0.150 to 0.271, $P = .57$) in short versus long children. The short children actually gained both less FFM ($-230 g$, 95% CI: -355 to -106 , $P < .001$) and FM ($-97 g$, 95% CI: -205 to 10), although the latter was not significant ($P = .076$). The differences in FM and FFM summed up to a 328 g (95% CI: 199 to 456, $P < .001$)—lower weight gain in short compared with long children. There were no differences in subscapular and triceps skinfold thickness and z scores (all $P > .5$).

For FM and FMI, there was no effect modification of product type (CSB or LNS) ($P > .30$). Likewise, there was no effect modification by stunting at admission (present or nonpresent) ($P > .50$) or stunting at admission stratified by severity (severe

TABLE 1 Baseline Characteristics of Children With MAM Included by MUAC Only With Complete D₂O Data (*n* = 390) by Length Category

	Short ^a	Long ^b	<i>P</i>
	<i>n</i> = 195	<i>n</i> = 195	
Girls, % (<i>n</i>)	84 (163)	72 (140)	.005
Age in mo, mean (±SD)	7.7 (1.6)	14.0 (4.3)	<.001
Wt in kg, mean (±SD)	6.012 (0.4)	7.487 (0.7)	<.001
Length in cm, mean (±SD)	63.9 (2.1)	72.0 (3.9)	<.001
MUAC in mm, mean (±SD)	121.0 (2.6)	122.2 (1.9)	<.001
WHZ, mean (±SD)	-1.5 (0.4)	-1.7 (0.3)	<.001
Wt-for-age z score, mean (±SD)	-2.4 (0.6)	-2.1 (0.6)	<.001
LAZ, mean (±SD)	-2.0 (1.0)	-1.7 (1.0)	.002
LAZ <-2, % (<i>n</i>)	49 (95)	35 (68)	.006
Season at time of inclusion, % (<i>n</i>)			.17
Dry season	61 (118)	67 (131)	
Rainy season	40 (77)	33 (64)	
Food supplement, % (<i>n</i>)			.84
CSB	48 (94)	47 (92)	
LNS	52 (101)	53 (103)	
Site, % (<i>n</i>)			.009
0	12 (24)	27 (52)	
1	15 (30)	11 (21)	
2	21 (41)	18 (35)	
3	31 (60)	26 (50)	
4	21 (40)	19 (37)	
Breastfeeding, % (<i>n</i>)	100 (194)	91 (178)	<.001

^a Short = <67 cm.

^b Long = ≥67 cm.

TABLE 2 Body Composition Among Short and Long Children With MAM at Baseline and Endline, Without Adjustments

	Baseline			Endline		<i>P</i> ^d Δ
	Short ^a	Long ^b	<i>P</i> ^c Δ	Short	Long	
	<i>n</i> = 195	<i>n</i> = 195		<i>n</i> = 195	<i>n</i> = 195	
FM in kg, mean (±SD)	1.098 (0.4)	1.218 (0.4)	.001	1.123 (0.4)	1.246 (0.4)	.003
FFM in kg, mean (±SD)	4.914 (0.5)	6.268 (0.8)	<.001	5.673 (0.5)	7.096 (0.9)	<.001
Fat percentage (FM/wt × 100), mean (±SD)	18.3 (5.8)	16.4 (5.3)	<.001	16.3 (5.7)	15.0 (4.9)	.015
FMI in kg/m ² , mean (±SD)	2.689 (0.9)	2.364 (0.8)	<.001	2.499 (0.9)	2.259 (0.8)	.006
FFMI in kg/m ² , mean (±SD)	12.000 (0.9)	12.056 (0.9)	.53	12.640 (0.8)	12.723 (0.9)	.36
Triceps skinfold in mm, mean (±SD)	6.6 (1.1)	6.3 (1.0)	.001	6.8 (1.1)	6.8 (1.2)	.85
Triceps skinfold-for-age z score, mean (±SD)	-1.5 (0.8)	-1.3 (0.8)	.04	-1.1 (0.8)	-0.8 (0.9)	.007
Subscapular skinfold in mm, mean (±SD)	5.6 (0.9)	5.1 (0.8)	<.001	5.8 (1.0)	5.4 (0.8)	<.001
Subscapular skinfold-for-age z score, mean (±SD)	-1.3 (0.9)	-1.5 (0.9)	.03	-0.9 (1.0)	-0.9 (0.9)	.96

^a Short = <67 cm.

^b Long = ≥67 cm.

^c *P* value for difference at baseline between short and long children.

^d *P* value for difference at endline between short and long children.

stunting, moderate stunting, or non-present) (*P* > .45).

The data set used is available from the corresponding author on reasonable request.

DISCUSSION

In response to WHO's call for research to identify appropriate MUAC admission and discharge criteria for children <67 cm and ≥6

months for supplementary foods,² we show here that short children with MUACs 115 to 125 mm but WHZs >-2 do not gain excessive fat during supplementation. Our results, therefore, can be used to challenge current policy and practice in several African countries of measuring MUAC only in children >6 months of age above a certain length threshold, thus excluding shorter children with only low MUACs from eligibility for MAM or SAM treatment programs.⁵⁻¹⁴

The exclusion of short children is based solely on expert opinion and reflects a concern that these children are just stunted rather than wasted, have slow catch-up growth, and may accumulate excessive FM after supplementation, thereby putting them at risk for noncommunicable diseases later in life. We previously reported that ponderal growth rates were similar in short and long children who received supplementation for MAM determined only by low MUAC,²⁰ and a similar finding was reported in children with SAM.²⁸

With this current study, we are the first to report direct evidence on fat accumulation after treatment of such children, and our data reveal there is no excessive fat gain in the short children compared with long children. In fact, weight gain in both groups overwhelmingly came in the form of FFM. These findings were independent of children having received LNS or CSB supplement. Because LNS contains a high level of fat, some experts were concerned that this food could lead to higher fat deposition in the malnourished child. Earlier we showed that LNS did not lead to high fat accumulation in the treatment of children for MAM.²² We now show that this concern is unwarranted even in shorter children enrolled in treatment solely by MUAC and,

TABLE 3 Changes in Outcomes During Supplementation

	Short ^a	Long ^b	Difference in Change ^c	P ^d Δ
	n = 195	n = 195		
FM in kg, mean (95% CI)	-0.022 (-0.088 to 0.043)	0.075 (0.010 to 0.140)	-0.097 (-0.205 to 0.010)	.076
FFM in kg, mean (95% CI)	0.678 (0.604 to 0.752)	0.908 (0.834 to 0.982)	-0.230 (-0.355 to -0.106)	<.001
Fat percentage (fat-mass/wt × 100), mean (95% CI)	-1.8 (-2.6 to -1.0)	-1.5 (-2.3 to -0.7)	-0.3 (-1.6 to 1.1)	.70
FMI in kg/m ² , mean (95% CI)	-0.166 (-0.309 to -0.234)	-0.128 (-0.271 to 0.015)	-0.038 (-0.257 to 0.181)	.74
FFMI in kg/m ² , mean (95% CI)	0.684 (0.556 to 0.813)	0.624 (0.495 to 0.752)	0.061 (-0.150 to 0.271)	.57
Triceps skinfold in mm, mean (95% CI)	0.39 (0.04 to 0.73)	0.35 (-0.002 to 0.70)	0.04 (-0.24 to 0.32)	.77
Triceps skinfold-for-age z score, mean (95% CI)	0.48 (0.25 to 0.71)	0.50 (0.26 to 0.73)	-0.02 (-0.22 to 1.85)	.88
Subscapular skinfold in mm, mean (95% CI)	0.32 (0.17 to 0.47)	0.24 (0.09 to 0.40)	0.08 (-0.15 to 0.30)	.52
Subscapular skinfold-for-age z score, mean (95% CI)	0.52 (0.36 to 0.67)	0.49 (0.34 to 0.65)	0.02 (-0.21 to 0.26)	.84

Data are mean and 95% CI calculated by using linear mixed models adjusted for food intervention, baseline measure of the outcome, age, sex, mo of admission, and site (random effects).

^a Short = <67 cm.

^b Long = ≥67 cm.

^c Difference in change of short versus long children.

^d P value for difference in change from baseline to endline for short and long children.

furthermore, is not modified by stunting at admission.

Using the state-of-the-art D₂O dilution technique seconded by skinfold thickness measurements, we present comprehensive data on body fat after supplementation in MAM. Because the gold standard for body composition is cadaver dissection, all in vivo techniques are necessarily imperfect. Here, we used 2 different techniques in which their error is uncorrelated, but their results are similar, making our findings more robust.

At baseline, the short children compared with the long children were younger, more often girls, had lower MUACs, and had higher prevalence of stunting and underweight but were less wasted.²⁰ More than 90% of the 1.5 kg–higher weight in long children at baseline was FFM, whereas short children had higher baseline FMI and fat percentage, and this was supported

by greater baseline skinfold thicknesses.

One possible explanation for this baseline difference is that, in the process of growing poorly, short children invested relatively more in fat reserves than long children, at a cost to their functional fat-free tissue. However, a simpler alternative explanation is that even in children with optimal growth conditions, such as those used to establish WHO 2006 growth standards, both triceps and subscapular skinfold thicknesses tend to decrease between 6 months and 2 years and more so in boys than girls. Hence, the fact that the short children were also ~6 months younger at baseline might explain the higher FMI at baseline. Supporting this, baseline skinfold thicknesses were thicker in short children, but when expressed as z scores relative to WHO reference data, short children actually had lower triceps values than long

children, whereas their subscapular skinfold thickness was only greater by a z score of 0.2 compared with long children.

At the end of the intervention, the average skinfold thicknesses in this study were less than a z score of -1 in both groups, suggesting no excessive fat gain in either group.

At the end of the supplementation, relative measures of FMI, FFMI, and fat percentage were similar between the 2 groups, and likewise there was no difference between groups in skinfold thickness changes, whether expressed in absolute terms or as z scores. The short children gained less weight and FFM and also showed a trend toward less FM gain.

In a recent study, it was found that children with MAM diagnosed solely on low MUAC experienced high rates of deterioration to SAM if left untreated.²⁹ We are not aware of data on mortality specifically for short children with MAM based only on low MUAC. However, the low-MUAC short children in our cohort exhibited a high prevalence of additional anthropometric deficits. Half of the short children were stunted, and short children were more underweight compared with long children. Stunted and underweight children have a higher mortality risk and may, therefore, benefit most from nutritional interventions.³⁰ MUAC is a simple way to target them for treatment.

WHO recently updated its Integrated Management of Childhood Illness and now recommends against routinely supplementing children with MAM because of a concern for obesity. We previously showed that there was no excess fat accumulation in children with MAM supplemented with LNSs or CSBs.²² In our findings here, we show that even the shorter children with MAM

determined solely by MUAC, who were considered by some experts to be at the greatest risk for obesity from food supplementation, do not gain excessive fat. Such children should no longer be excluded from treatment eligibility.

We are not aware of any data on fat accumulation after treatment in short children with SAM by MUAC alone. Indeed, currently only data on skinfolds could be generated in large field studies on children with SAM because current techniques available for body composition assessment are not suitable for use in SAM. However, we see no rationale in maintaining a length restriction for MUAC assessment in SAM.

CONCLUSIONS

Short children with low MUAC do not gain excessive fat during supplementation. Admission by the same MUAC criteria (ie, <115 mm for SAM and 115–125 mm for MAM) should apply to any child 6 to 59 months of age, regardless of their length and irrespective of their WHZ. The use of length as criterion for measuring MUAC should be discontinued in policy and practice wherever such restrictions exist. It will also be important for WHO to integrate this evidence, in which we show that children with MAM are not put at risk for obesity after food supplementation, into its recently updated Integrated Management of Childhood Illness recommendations.

ABBREVIATIONS

CI: confidence interval
CSB: corn-soy blend
D₂O: deuterium oxide
FFM: fat-free mass
FFMI: fat-free mass index
FM: fat mass
FMI: fat mass index
LAZ: length-for-age z score
LNS: lipid-based nutrient supplement
MAM: moderate acute malnutrition
MUAC: mid-upper arm circumference
SAM: severe acute malnutrition
TBW: total body water
WHO: World Health Organization
WHZ: weight-for-height z score

to data interpretation; Drs Michaelsen and Friis conceptualized the research and design and contributed to the interpretation of data; and all authors critically revised the article for important intellectual content, approved the final manuscript as submitted, and agree to be accountable for all aspects of the work.

This trial has been registered at www.controlled-trials.com (identifier ISRCTN42569496).

DOI: <https://doi.org/10.1542/peds.2018-0679>

Accepted for publication Jun 6, 2018

Address correspondence to Christian Fabiansen, MD, PhD, Department of Nutrition, Sports and Exercise, University of Copenhagen, Rolighedsvej 25, Frederiksberg, 1958 Copenhagen, Denmark. E-mail: chfa@nexs.ku.dk

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Other than those listed in Potential Conflict of Interest, the authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Funded by Danish International Development Assistance (grant 09-097LIFE) (K.F.M.); Médecins Sans Frontières (Denmark, Norway); Arvid Nilsson's Foundation; The World Food Program, which was part of a donation to the World Food Program from the American people through the support of the US Agency for International Development's Office of Food for Peace; the Alliance for International Medical Action; and the European Union's humanitarian aid funds, in partnership with Action Contre la Faim. This article covers humanitarian aid activities implemented with the financial assistance of the European Union. The views expressed herein should not be taken, in any way, to reflect the official opinion of the European Union, and the European Commission is not responsible for any use that can be made of the information it contains. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the article.

POTENTIAL CONFLICT OF INTEREST: Dr Michaelsen has received research grants from the US Dairy Export Council and the Danish Dairy Research Foundation and also has research collaboration with Nutriset, the main producer of lipid-based nutrient supplement products; Mr Phelan serves on Nutriset's Social Purposes Advisory Commission; Dr Briend was the inventor of lipid-based nutrient supplements but abandoned claims to royalties in 2003; Dr Friis has received research grants from Arla Food for Health Centre and also has research collaboration with Nutriset; the other authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. World Health Organization. Children: reducing mortality. Available at: www.who.int/mediacentre/factsheets/fs178/en/. Accessed June 25, 2018
2. World Health Organization. *UNICEF, WFP and UNHCR Consultation on the Programmatic Aspects of the Management of Moderate Acute Malnutrition in Children Under Five Years of Age*. Geneva, Switzerland: World Health Organization; 2010
3. Black RE, Victora CG, Walker SP, et al; Maternal and Child Nutrition Study Group. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382(9890):427–451
4. Olofin I, McDonald GM, Ezzati M, et al; Nutrition Impact Model Study (Anthropometry Cohort Pooling). Associations of suboptimal growth with all-cause and cause-specific

- mortality in children under five years: a pooled analysis of ten prospective studies. *PLoS One*. 2013;8(5):e64636
5. Cameroon Ministry of Public Health. *National Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2013;20, 29, 119
 6. Central African Republic Ministry of Public Health and Population. *Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2014;40, 57, 168
 7. Republic of Chad Ministry of Public Health, Social Action and National Solidarity. *Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2014:166
 8. Guinea Ministry of Health and Public Hygiene. *National Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2012:36, 52, 149
 9. Ivory Coast Ministry of Public Health and Hygiene. *National Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2015:56, 83, 188
 10. Islamic Republic of Mauritania Ministry of Health. *Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2011:57, 146
 11. Republic of Senegal Ministry of Health and Social Action. *Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2013:26, 40, 57, 156
 12. Togo Ministry of Health. *National Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2013:33, 123
 13. Mali Ministry of Health. *Protocol for the Integrated Management of Acute Malnutrition in Mali [in French]*. 2011:31, 138
 14. Ethiopia Federal Ministry of Health. *Protocol for the Management of Severe Acute Malnutrition*. 2007:10
 15. World Health Organization. *The Management of Nutrition in Major Emergencies*. Geneva, Switzerland: WHO; 2000
 16. World Health Organization, Regional Office for the Eastern Mediterranean. *Field Guide on Rapid Nutritional Assessment in Emergencies*. Alexandria, Egypt: WHO; 1995
 17. Médecins Sans Frontières. *MSF Nutrition Guidelines*. 1995. Available at: <http://www.nzdl.org/gsdmod?e=d-00000-00---off-0cdl--00-0---0-10-0--0--0direct-10--4-----0-11--11-en-50--20-about--00-0-1-00-0-4---0-0-11-10-OutfZz-8-00&cl=CL1.155&d=HASH9237a20340ca39bd07dda2&gc=1>. Accessed July 16, 2018
 18. Valid International. *Community-Based Therapeutic Care (CTC): A Field Manual*. Oxford, United Kingdom: Valid International; 2006
 19. World Health Organization. Assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition. Available at: www.who.int/nutrition/publications/guidelines/children-primaryhealthcare-obesity-dbm/en/. Accessed November 1, 2017
 20. Fabiansen C, Phelan KP, Cichon B, et al. Short children with a low midupper arm circumference respond to food supplementation: an observational study from Burkina Faso. *Am J Clin Nutr*. 2016;103(2):415–421
 21. Niger Ministry of Public Health. *Protocol for the Integrated Management of Acute Malnutrition [in French]*. 2016:127
 22. Fabiansen C, Yaméogo CW, Iuel-Brockdorf A-S, et al. Effectiveness of food supplements in increasing fat-free tissue accretion in children with moderate acute malnutrition: a randomised 2 × 2 × 3 factorial trial in Burkina Faso. *PLoS Med*. 2017;14(9):e1002387
 23. Fabiansen C, Yaméogo CW, Devi S, Friis H, Kurpad A, Wells JC. Deuterium dilution technique for body composition assessment: resolving methodological issues in children with moderate acute malnutrition. *Isotopes Environ Health Stud*. 2017;53(4):344–355
 24. International Atomic Energy Agency. *Introduction to Body Composition Assessment Using the Deuterium Dilution Technique With Analysis of Saliva Samples by Fourier Transform Infrared Spectrometry*. Vienna, Austria: International Atomic Energy Agency; 2011
 25. Davies PS, Wells JC. Calculation of total body water in infancy. *Eur J Clin Nutr*. 1994;48(7):490–495
 26. Racette SB, Schoeller DA, Luke AH, Shay K, Hnilicka J, Kushner RF. Relative dilution spaces of 2H- and 18O-labeled water in humans. *Am J Physiol*. 1994;267(4, pt 1):E585–E590
 27. Lohman TG, ed. Estimating body composition in children and the elderly. In: *Advances in Body Composition Assessment (Current Issues in Exercise Science, Monograph 3)*. Champaign, IL: Human Kinetics; 1992:65–77
 28. Dale NM, Myatt M, Prudhon C, Briend A. Using mid-upper arm circumference to end treatment of severe acute malnutrition leads to higher weight gains in the most malnourished children. *PLoS One*. 2013;8(2):e55404
 29. James P, Sadler K, Wondafrash M, et al. Children with moderate acute malnutrition with no access to supplementary feeding programmes experience high rates of deterioration and no improvement: results from a prospective cohort study in rural Ethiopia. *PLoS One*. 2016;11(4):e0153530
 30. McDonald CM, Olofin I, Flaxman S, et al; Nutrition Impact Model Study. The effect of multiple anthropometric deficits on child mortality: meta-analysis of individual data in 10 prospective studies from developing countries. *Am J Clin Nutr*. 2013;97(4):896–901

**Short Malnourished Children and Fat Accumulation With Food
Supplementation**

Christian Fabiansen, Kevin P.Q. Phelan, Bernardette Cichon, Charles W. Yaméogo,
Ann-Sophie Iuel-Brockdorff, Anura Kurpad, Jonathan C. Wells, Christian Ritz,
Suzanne Filteau, André Briend, Vibeke B. Christensen, Kim F. Michaelsen, Susan
Shepherd and Henrik Friis

Pediatrics originally published online August 24, 2018;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/early/2018/08/22/peds.2018-0679
References	This article cites 10 articles, 2 of which you can access for free at: http://pediatrics.aappublications.org/content/early/2018/08/22/peds.2018-0679#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Nutrition http://www.aappublications.org/cgi/collection/nutrition_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Short Malnourished Children and Fat Accumulation With Food Supplementation

Christian Fabiansen, Kevin P.Q. Phelan, Bernardette Cichon, Charles W. Yaméogo,
Ann-Sophie Iuel-Brockdorff, Anura Kurpad, Jonathan C. Wells, Christian Ritz,
Suzanne Filteau, André Briand, Vibeke B. Christensen, Kim F. Michaelsen, Susan
Shepherd and Henrik Friis

Pediatrics originally published online August 24, 2018;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2018/08/22/peds.2018-0679>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

